

Visualisation of vascular cannabinoid receptors and their potential interaction with α_1 -adrenergic receptors.

Daly CJ, Wallace G, White K, Henstridge C*, Irving A* & McGrath JC.

Integrative & Systems Biology, FBLS, University of Glasgow. Glasgow G12 8QQ.

*Centre for Neuroscience, Division of Molecular and Translational Medicine, University of Dundee, Ninewells Hospital & Medical School, Dundee DD1 9SY

The role of cannabinoid receptors (CB1 & CB2) within the cardiovascular system is unclear. The endogenous cannabinoid anandamide (AEA) mediates vasodilation *in-vitro* [1] whilst *in-vivo* a triphasic blood pressure response comprising pressor and depressor components has been reported [2]. A possible role for both CB1 and CB2 receptors exists within vascular tissue. However, non-CB mediated responses in the vasculature have been observed and a role for the orphan receptor GPR55 has been postulated [3]. The aims of this study were to a) investigate the role of endogenous cannabinoids in mouse tail artery, a thermoregulatory vessel rich in α_1 -adrenoceptors and b) examine, for the first time, the binding of a novel fluorescent ligand for CB receptors (Tocrifluor 1117).

Tail arteries were removed from 4 month old male C57 black mice. Vessel segments were either mounted in a wire myograph for functional studies or incubated in Tocrifluor 1117 (0.5 μ M) & QAPB (1 μ M, fluorescent α_1 -adrenoceptor antagonist) for confocal analysis. Concentration response curves (CRC) to noradrenaline (NA) were performed in the presence and absence of the endocannabinoids AEA (1 μ M) and 2-arachidonylglycerol (2-AG, 1 μ M). Tocrifluor 1117 and QAPB were imaged under 543nm and 488nm excitation respectively. Tocrifluor 1117 binding was also examined in HEK293 cells stably expressing GPR55.

AEA 1 μ M caused a transient contraction when applied to isolated tail artery segments (0.08g). The NA CRC was shifted to left in the presence of 1 μ M AEA (Log EC₅₀ control -6.80 vs -7.82, $p < 0.05$). The maximum contractile response was unchanged. In the presence of 2-AG (1 μ M) a small leftward shift of the NA CRC was observed which failed to reach significance. However comparison of the effect of 2-AG alone (Log EC₅₀ -7.36) and 2-AG plus indomethacin (10 μ M, Log EC₅₀ -6.33) revealed a significant difference ($p > 0.05$). Tocrifluor 1117 binding was observed on all 3 vascular tunics with particularly strong fluorescence evident on perivascular fat, adventitial cells, nerves and smooth muscle. In several areas of media, colocalisation of Tocrifluor 1117 and QAPB was observed. In live HEK293 GPR55 cells, Tocrifluor 1117 generated a rise in Ca⁺⁺ and promoted receptor clustering, visible as punctate fluorescence which developed over time.

Tocrifluor 1117 (fluorescent analogue of the CB1 antagonist AM251) is a potentially very powerful tool for identifying the cellular location of cannabinoid receptors, including GPR55 in living tissues. AM251 has been shown to activate GPR55 [4]. The results of the study suggest that endogenous cannabinoids potentiate the actions of noradrenaline in tail artery possibly via co-localised cannabinoid and α_1 -adrenergic receptors in vascular smooth muscle. The importance of the fluorescent binding in perivascular fat and adventitia requires further study.

1. Hogestadt ED & Zygmunt PM (2002) Cardiovascular Pharmacology of Anandamide. Prostaglandins, Leukotrienes & Essential Fatty Acids. 66 (2&3); 343-351.
2. Varga K, et al., (1996) Mechanism of the hypotensive action of anandamide in anesthetized rats. Hypertension. Oct;28(4):682-6
3. Hiley CR & Kaup SS (2007) GPR55 and the vascular receptors for cannabinoids. Br. J. Pharmacol., 152, 559-561

4. Henstridge C et al., (2008). The GPR55 ligand L- α -lysophosphatidylinositol promotes RhoA-dependent Ca²⁺ signaling and NFAT activation. *FASEB J. Pub Online Aug 2008.*